

# **An Attribution Study of the UK mean temperature in summer 2026**

Georgie Logan, Ishbel Scott, Andrew Ciavarella, Mark  
McCarthy

## Document history

| Version | Purpose                   | Date       |
|---------|---------------------------|------------|
| 0.1     | Draft                     | 01-09-2026 |
| 1.0     | Revision following review | 07-09-2026 |

### Prepared by

Georgie Logan

Ishbel Scott

Andrew Ciavarella

Mark McCarthy

### Reviewed by

Nick Rayner, Strategic Head, Climate Monitoring and Attribution

### Authorised for issue by

Nick Rayner, Strategic Head, Climate Monitoring and Attribution

## Contents

To update the table of contents, right-click on it and select 'Update Field'.

|                             |    |
|-----------------------------|----|
| Document history .....      | 2  |
| Contents .....              | 3  |
| 1. Introduction .....       | 4  |
| 2. Data .....               | 6  |
| 3. Method.....              | 8  |
| 4. Model Evaluation .....   | 8  |
| 5. Climate Attribution..... | 11 |
| References .....            | 13 |

## 1. Introduction

Summer 2026 was the hottest on record for the UK. Persistently higher than average temperatures, with multiple heatwave events defined the summer season ([Figure 1](#)). Dominant high-pressure systems also led to the second sunniest summer for England and Wales, and the UK's third driest July. Heatwaves in both June and August saw peak maximum daily temperatures at or above 38°C. 38.0 °C at Lingwood (Norfolk) on 26 June set a new maximum daily temperature record for the month of June, and 38.1°C at Kew Gardens (Greater London) on 13 August was provisionally the highest temperature recorded during the season. The UK summer mean temperature of 16.5°C means 2026 replaces 2025 (16.1°C) as the hottest summer on record, [as shown in Figure 2](#).

A rapid attribution analysis was conducted to quantify the human-influence on the likelihood of reaching or exceeding the 2026 summer mean temperature. The approach provides estimates of the exceedance probability of similar events in the climate of a 'natural' world, the climate of the present day, and that of the end of the 21<sup>st</sup> Century.

The analysis was conducted using a system developed by the Met Office for rapid studies and described in Christidis (2021). This uses climate models from the Coupled Model Intercomparison Project Phase 6 (CMIP6, Eyring et al., 2016). This system assesses the likelihood of reaching (or exceeding) the UK summer mean temperature of 2026 in the current climate, and in one without human-induced climate change, to estimate how climate change, primarily through the emission of greenhouse gases, has changed the probability of such an event. It does not condition these probabilities on the specific large-scale circulation, and therefore does not separate out the role that specific regional or remote dynamical drivers may have played.




Figure 1: UK daily mean temperature for summer 2026. Periods of above average temperature are highlighted orange, and periods below average are blue. The 1991-2020 average temperature is shown in black. The (light) grey shaded regions show the (5%-95%) 10%-90% of all summers, and the thin orange and blue outer lines show the daily record temperatures. The thick blue line shows 2026 daily mean temperatures.




Figure 2: UK summer mean temperature from 1884 to 2026. The smoothed trend is shown as a black dashed curve. The 1991-2020 summer average is shown as a pink line.

## 2. Data

Observed values of the UK summer mean temperature are obtained from the HadUK-Grid dataset v1.3.2.0 (Hollis et al., 2019; Met Office, 2026). The time series contains UK monthly means spanning January 1884 – August 2026, with the 2026 values being provisional at the time of writing. Summer mean values are obtained by averaging this monthly data over summer months of June, July and August, appropriately weighted by the number of days in each month.

Model data is taken from 14 CMIP6 models that provide ensemble members for each of the natural (hist-nat) and historical (historical) climate experiments as well as projections from the SSP2-4.5 scenario (ssp245, Riahi et al., 2017). The hist-nat experiment provides simulations of the pre-industrial climate and the historical experiment provides simulations of the changing climate under observed climate forcings. The SSP2-4.5 projections are valid from 2015 onwards and are chosen as a “middle of the road” scenario for mitigation and adaptation to climate change. These projections see moderate global population growth and uneven development, that may remain a close analogue of the real world. The names of models and the number of ensemble members used from each are shown in Table 1.

Table 1. CMIP6 models and corresponding number of ensemble members used from each experiment in the attribution analysis.

| Model Name             | historical | ssp245    | hist-nat  |
|------------------------|------------|-----------|-----------|
| <b>ACCESS-CM2</b>      | 3          | 3         | 3         |
| <b>ACCESS-ESM1-5</b>   | 18         | 18        | 3         |
| <b>BCC-CSM2-MR</b>     | 3          | 1         | 3         |
| <b>CESM2</b>           | 11         | 6         | 3         |
| <b>CNRM-CM6-1</b>      | 30         | 6         | 10        |
| <b>CanESM5</b>         | 25         | 25        | 15        |
| <b>FGOALS-g3</b>       | 4          | 4         | 3         |
| <b>GFDL-ESM4</b>       | 3          | 3         | 3         |
| <b>GISS-E2-1-G</b>     | 6          | 3         | 5         |
| <b>HadGEM3-GC31-LL</b> | 5          | 5         | 5         |
| <b>IPSL-CM6A-LR</b>    | 32         | 11        | 10        |
| <b>MIROC6</b>          | 10         | 3         | 10        |
| <b>MRI-ESM2-0</b>      | 5          | 1         | 5         |
| <b>NorESM2-LM</b>      | 3          | 3         | 3         |
| <b>Total</b>           | <b>158</b> | <b>92</b> | <b>81</b> |

Time series of the observed and modelled UK summer mean temperature anomalies are shown in Fig. 3. Anomalies are calculated from a baseline of average summer temperature for the 30-year period 1901 to 1930. Increasing summer temperatures are shown in both observations and the historical and ssp245 simulations, with further increases projected through the rest of the 21<sup>st</sup> Century. This shows a steadily increasing likelihood of

significantly higher-than-baseline summer temperatures being experienced in the UK. The historical and hist-nat ensembles start to diverge from around the 1980s suggesting that climate forcings arising from human influence are the dominant factor in the long-term observed warming trend.

The value of the UK summer mean temperature anomaly in 2026, relative to a baseline period of 1901 – 1930, was  $+3.01^{\circ}\text{C}$ ,  $0.4^{\circ}\text{C}$  above the record set in 2025 of  $+2.61^{\circ}\text{C}$ . 2026 is estimated as a relatively hot summer for the current climate, but higher extremes are still possible.




Figure 3: Time series of the UK summer mean temperature anomaly (w.r.t. 1901 – 1930). Observational data from HadUK-Grid (black). Simulations from the CMIP6 historical (red), ssp245 (purple) and hist-nat (green) experiments. Simulation data is represented as median values and filled 5<sup>th</sup> – 95<sup>th</sup> percentile ranges explored by members of the multi-model ensemble. Simulation data is smoothed with a rolling window of 20 years, with historical and ssp245 combined into one continuous series. Observed data runs from 1884 – 2026.

### 3. Method

The method employed is to find exceedance probabilities of the 2026 UK summer mean temperature in the natural world climate without human influence (*nat*), the current day climate (*now*) and that of the end of the century (*end*). The *nat* climate is represented by collating all available years from the hist-nat simulations (1850 - 2020). The *now* climate is represented by taking all data from the 20-year period 2016 – 2035 that is centred on 2026, from the historical simulations up to 2014 and continued by ssp245. The *end* climate is represented by all data taken from the final 20 years of the ssp245 experiment (2081 - 2100).

Time series are produced from both observations and model members of UK regional mean anomalies of gridded summer mean temperatures. In all cases the anomalies are calculated relative to a 1901 – 1930 baseline. Land fraction masks are applied for each model individually to obtain land-only temperatures comparable to the HadUK-Grid observations. All available members from the 14 CMIP6 models are combined into a single multi-model ensemble and samples taken from the appropriate time periods, from which distributions are estimated.

Exceedance probabilities are then calculated in one of two ways.

If the observed value of the UK summer mean temperature anomaly is not an extreme value, then the exceedance probability is the empirical fraction of model values lying above the observed threshold. An extreme value is one that sits within the tail of the model data distribution, which we define as the top 20% of values.

If the observed value of the UK summer mean temperature anomaly can be treated as an extreme value, then a Generalised Pareto Distribution (GPD) is fit to the model tail data and an exceedance probability is found from the continuous approximation of the tail of the model distribution.

Probability ratios are used to express changes in probability between the current climate and that of the natural world, and between the climate of the end of the century and the natural world. The probability ratio in each case is the exceedance probability in the climate of interest divided by that in the pre-industrial climate.

Uncertainty ranges are calculated by performing a 10,000-member bootstrap (90% with replacement) from which the 5% and 95% values of the exceedance probabilities, return times and probability ratios are found. For each bootstrap resampling of the data, we re-count or re-fit the GPD to find the new exceedance probability.

### 4. Model Evaluation

The models are evaluated against the HadUK-Grid observations, as presented in Figure 4 and Figure 5. Several evaluation tests commonly employed in event attribution studies (Christidis et al., 2013, Christidis, 2021) are conducted on the full multi-model ensemble of



historical simulations over the evaluation period 1884 – 2014, which is the overlap of the historical experiment with availability of the observations (see Figure 4). This provides over 20,000 years of model data. In Figure 5, the same validation tests are also conducted on “recent” model and observation data, defined as the last 50 years (1977 to 2026), where recent model data are provided by the relevant years of the historical and SSP2-4.5 experiments combined. This gives us confidence that the observational data has continued to be represented well by the model ensemble in the time periods we are now evaluating.

The observed temperature trend is well within the range of the simulations and very close to the ensemble mean (Figure 4 and Figure 5, top panel, observed and model mean trend indistinguishable in Figure 4). A small number of simulations exhibit a negative trend over the evaluation period, but we do not exclude these members. Examining each model separately, all models have a positive trend when extended further into the 21st Century with the SSP2-4.5 scenario, which is clear in the recent validation trend plot in Figure 5.

Estimates of the power spectra as seen in the periodogram (Figure 4 and Figure 5, middle panel) also indicate good consistency of the variability in summer mean temperature between the model ensemble and HadUK-Grid, both at short and long periods.

The Quantile-Quantile plot (Figure 4 and Figure 5, lower panel) produced for each simulation separately shows curves that mostly lie close to the 1:1 line, indicating that the shapes of member distributions compare well with that observed. The presence of model values cooler than observed below anomalies of 0°C in Figure 4 expresses the fact that the observed world explored cooler variations during the anomaly period 1901 – 1930 (Figure 3). Model members examined separately exhibit consistent variability at multi-decadal time scales (see the periodograms, Figure 4 and Figure 5, middle panel). As the observed quantiles still remain within the range of members over these values, and both trend and periodogram show consistency, we accept the multi-model ensemble representation of the shape of the distribution of anomalies.

We do not exclude any members or models based on these assessments. The ensemble is deemed suitable for an attribution analysis of extreme UK summer mean temperatures.




Figure 4: Evaluation of CMIP6 multi-model ensemble used in the attribution analysis, for the historical experiment, evaluated over 1884 – 2014 by comparison with observational HadUK-Grid data. Top panel: temperature trends over the evaluation period. Middle panel: power spectra estimates (periodogram) from the HadUK-Grid data and historical simulations. Bottom panel: quantile-quantile plot for each of the historical simulations.




Figure 5: Evaluation of CMIP6 multi-model ensemble used in the attribution analysis, for recent experiments (combined historical and ssp245 experiments), evaluated over 1977 – 2026 by comparison with observational HadUK-Grid data. Top panel: temperature trends over the evaluation period. Middle panel: power spectra estimates (periodogram) from the HadUK-Grid data and recent simulations. Bottom panel: quantile-quantile plot for each of the recent simulations.

## 5. Climate Attribution

The results of the analysis are presented as the probability and equivalent return period of exceeding the 2026 observed summer mean temperature anomaly, along with associated probability ratios in Table 2. UK summer mean temperatures exceeding those observed in 2026 are expected to occur with a best estimate return time of approximately 9 years (5% to 95% range 8.2 to 9.2 years) in the current climate, human influence having increased the probability of occurrence by around 130 times from an equivalent return time estimate of 1148 years (5% to 95% range 780 to 1800 years) in the climate of the pre-industrial period. There is more uncertainty in the pre-industrial period's return time as the threshold is notably more extreme for the natural model data.

Individual model projections (not shown) diverge by the end of the century (apparent from Figure 3) but further warming to the end of the century could see the temperatures of 2026 being exceeded as frequently as every other year. The multi-model best estimate return time is approximately 1.5 years at the end of the century, but this includes a range of projections and is shorter in some individual model projections and longer in others.

Table 2. Exceedance probabilities and risk ratio estimates for exceeding the observed 2026 UK summer mean temperature anomaly (w.r.t. 1901 - 1930). 5-95% uncertainty ranges found by a 10,000-member bootstrap.

| Experiment                                          | Exceedance Probability<br>(5%, 95%) | Return time / years<br>(5%, 95) | Probability Ratio<br>(5%, 95%) |
|-----------------------------------------------------|-------------------------------------|---------------------------------|--------------------------------|
| <b>nat: hist-nat<br/>all data</b>                   | 0.0008711 (0.00057, 0.0013)         | 1148 (780, 1800)                | N/A                            |
| <b>now: historical<br/>and ssp245<br/>2016-2035</b> | 0.1152 (0.11, 0.12)                 | 8.681 (8.2, 9.2)                | 132.2 (89, 200)                |
| <b>end: ssp245<br/>2081-2100</b>                    | 0.6543 (0.63, 0.67)                 | 1.528 (1.5, 1.6)                | 751.2 (510, 1200)              |

Other attribution studies have evaluated UK summer mean temperature extremes using essentially the same method as used here. McCarthy et al. (2019) conducted an attribution study of the, at the time, record 2018 summer. This study estimated a return time of reaching or exceeding 2018 summer mean temperature of 8 to 9 years, for a climate centred on 2018. In Kendon et al. (2024) a corresponding analysis but for just England, rather than the UK, and for the summer of 1976 found a return time in the current climate of 4.25 years.

In last year's version of this report, Logan et al. (2025) stated that the 2025 summer record was "not particularly extreme in the current climate, and that considerably higher summer mean temperatures are already possible". 2026 has demonstrated that this summer.

## References

Christidis, N., P. A. Stott, A. Scaife, A. Arribas, G. S. Jones, D. Copsey, J. R. Knight, and W. J. Tennant, 2013: A new HadGEM3-A based system for attribution of weather and climate-related extreme events, *J. Clim.*, 26, 2756-2783, <https://doi.org/10.1175/JCLI-D-12-00169.1>.

Christidis, N., 2021: Using CMIP6 multi-model ensembles for near real-time attribution of extreme events, Hadley Centre Technical Note 107, Available from

<https://www.metoffice.gov.uk/research/library-and-archive/publications/science/climatescience-technical-notes>

Eyring, V., S. Bony, G. A. Meehl, C. A. Senior, B. Stevens, R. J. Stouffer, and K. E. Taylor, 2016: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci. Mod. Dev.*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>.

Hollis, D., M. McCarthy, M. Kendon, T. Legg, and I. Simpson, 2019: HadUK-Grid—A new UK dataset of gridded climate observations, *Geosci. Data. J.*, 6, 151-159, <https://doi.org/10.1002/gdj3.78>.

Kendon, E.J., Ciavarella, A., McCarthy, M., Brown, S., Christidis, N., Kay, G., et al. (2024) Multiperspective view of the 1976 drought–heatwave event and its changing likelihood. *Quarterly Journal of the Royal Meteorological Society*, 150(758), 232–261. <https://doi.org/10.1002/qj.4594>

McCarthy, M., Christidis, N., Dunstone, N., Fereday, D., Kay, G., Klein-Tank, A., Lowe, J., Petch, J., Scaife, A. and Stott, P. (2019), Drivers of the UK summer heatwave of 2018. *Weather*, 74: 390-396. <https://doi.org/10.1002/wea.3628>

Met Office; Hollis, D.; Carlisle, E.; Kendon, M.; Packman, S.; Doherty, A. (2025): HadUK-Grid Gridded Climate Observations on a 1km grid over the UK, v1.3.1.ceda (1836-2024). NERC EDS Centre for Environmental Data Analysis, 01 July 2025. doi:10.5285/f02cc6ddd92f45b18b9ab6ab544df7d9. <https://dx.doi.org/10.5285/f02cc6ddd92f45b18b9ab6ab544df7d9>

Riahi, K., and co-Authors, 2017: The shared socioeconomic pathways and their energy, land use, and greenhouse gas emissions implications: an overview, *Global Env. Change*, 42, 153– 168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>.

Met Office; Logan, G.; Ciavarella, A.; McCarthy, M.; (2025): An Attribution Study of the UK mean temperature in summer 2025, [https://www.metoffice.gov.uk/api/assets/file/hctn\\_summer\\_2025\\_analysis\\_v11pdf?prefix=assets](https://www.metoffice.gov.uk/api/assets/file/hctn_summer_2025_analysis_v11pdf?prefix=assets)